

Published on *British Columbia Drug and Poison Information Centre (BC DPIC)* (http://www.dpic.org)

Home > Printer-friendly PDF > Printer-friendly PDF

May is Food Allergy Awareness Month

Access: professional

Article type:

drug information

Food allergies appear to be on the rise in industrialised countries. They have been dubbed "the second wave of the allergy epidemic". Food allergies lessen quality of life and can be life-threatening. 2,3

Food Allergy Vs Intolerance

Food allergies are due to an immune response to food or food components and may be IgE-or non-IgE-mediated.^{3,4} Food allergies can present as anaphylaxis. However, food allergy can present with a wide range of signs and symptoms involving the gastrointestinal, cutaneous, and respiratory systems or a combination. Food intolerances, on the other hand, are mainly non-immune mediated reactions and include lactose intolerance, glutamate and aspartame sensitivity, or reactions to preformed toxins or vasoactive amines in foods.

Food Allergies And Medications

There are more than 170 foods known to cause IgE reactions,⁴ but most food allergies are caused by peanuts, tree nuts, seafood (fish and shellfish), eggs, milk, sesame, soy, and wheat.⁵ Medications may contain some of these potential allergens (Table 1). Some medications can worsen allergic reactions to food (Table 2).

Table 1: Potential food allergens found in some medications*

Peanut oil, soybean oil, sesame oil (in capsules, depot injections, lipid emulsions, nasal and topical products)	Refined peanut and soybean oils have low allergen content a suggest that ingestion is safe, but since there is no threshold known allergy, avoidance is still recommended. Data on administration is lacking and severe allergy to lipid emulsion reported. Most sesame oils are not refined enough to be safe
Soy lecithin (in capsules, tablets, oral liquids) and egg lecithin (in lipid emulsions)	Although some soy-allergic patients can eat soy lecithin, and seems to be safe in cooked foods, intravenous administration serious reactions. 10
Egg protein (in vaccines)	MMR vaccine and Imovax® rabies vaccine are generally safe patients. 11 Influenza vaccine is not contraindicated in egg all administration with trained resuscitation personnel and equip recommended. 12 Avoid yellow fever and RabAvert® rabies v
Wheat	Wheat allergy is different from celiac disease. 13 Most medicing free. Wheat starch is found in some tablets (e.g. Imunovir®, Tardan® shampoo contains hydrolyzed wheat protein.
Seafood	Protamine (NPH insulin, heparin antidote) is isolated from sa may be contaminated with fish proteins. Patients with fish alle lodine-containing drugs like amiodarone, contrast dyes, or pare not contraindicated in patients with fish or shellfish allergy is a misnomer. 15
Tree nuts	Medications do not contain tree nuts, but paclitaxel is found in patient with hazelnut allergy developed hives, itching and dysafter a paclitaxel infusion was started. 16
Miscellaneous (non-immune reactions)	Salicylates are found in fruits, vegetables, herbs and spices to by most patients with ASA sensitivity. To Carmine red was the anaphylaxis purported to be due to azithromycin. Sulfite se same as sulphonamide allergy.

^{*}Not a comprehensive list. Consult specific product information.

Table 2: Medications that can exacerbate food allergies

	May predispose to bronchoconstriction. ¹⁹ ASA increases foo exercise-induced anaphylaxis (possibly by increasing GI upta 20

	Do not worsen allergies <i>per se</i> but may blunt the therapeutic epinephrine. ³
ACE inhibitors	May interfere with production of endogenous vasoconstrictors allergic reactions. ³
Tacrolimus	Tacrolimus immunosuppression is associated with increased

Prevention Of Food Allergies

For patients with known food allergies avoiding the allergen is the goal. Thus, if a patient is suspected a having a food allergy, they must have the diagnosis confirmed and the specific allergen determined. Probiotics have been shown to reduce eczema, but there is insufficient evidence that probiotics prevent or treat food allergies.^{22,23}

Treatment Of Anaphylaxis

Details on treatment are beyond the scope of this article - see Additional Resources.

First-line: epinephrine. Guidelines and local experts recommend a 0.3 mg dose for children weighing more than 20 to 25 kg.^{3,24} Optimally, patients should have a back-up dose available, and all patients with a severe reaction should be sent immediately to the hospital.²⁵

Accidental injection of epinephrine into an extremity such as a finger can cause tissue ischemia. However, such accidents can sometimes be managed at home - contact the Poison Control Centre immediately.

Adjunctive treatments: Bronchodilators can help relieve bronchospasm not responding to epinephrine, but they do not relieve laryngeal edema and **EPINEPHRINE MUST BE GIVEN**. Antihistamines relieve itching and urticaria but do not relieve shortness of breath, wheezing or shock and **EPINEPHRINE MUST BE GIVEN**. Corticosteroids take several hours to work and are not useful for immediate treatment and **EPINEPHRINE MUST BE GIVEN**.³

Additional Resources

 Learn more about food allergies and anaphylaxis, and print out a personalised emergency treatment plan at www.allergysafecommunities.ca

- Health Canada food allergies and intolerances: http://www.hc-sc.gc.ca/fn-an/securit/allerg/index-eng.php
- Guidelines for the diagnosis and management of food allergy in the United States (2010) http://www.niaid.nih.gov/topics/foodallergy
- Patients can call 811 and speak to a HealthLinkBC allergy dietician.

Written by Raymond Li, BSc(Pharm), MSc. Reviewed by Laird Birmingham, MD, FRCPC, MHSc

Thank you to Linda Kirste, RD, Dietician Services, HealthLinkBC, for her expert advice.

References

- 1. Prescott S, Allen KJ. Food allergy: riding the second wave of the allergy epidemic. Pediatr Allergy Immunol. 2011; 22: 155-60.
- 2. Ravid NL, et al. Mental health and quality-of-life concerns related to the burden of food allergy. Immunol Allergy Clin N Am. 2012; 32: 83-95.
- 3. Boyce JA, et al. Guidelines for the diagnosis and management of food allergy in the United States: Report of the NIAID-Sponsored Expert Panel. J Allergy Clin Immunol. 2010; 126(6 Suppl): S1-S58.
- 4. Burks AW, et al. ICON: food allergy. J Allergy Clin Immunol. 2012 (in press). doi:10.1016/j.jaci.2012.02.001
- 5. Allen KJ, Koplin JJ. The epidemiology of IgE-mediated food allergy and anaphylaxis. Immunol Allergy Clin N Am. 2012; 32: 35-50.
- 7. Madsen CB, et al. Can we define a tolerable level of risk in food allergy? Report from a EuroPrevall/UK Food Standards Agency workshop. Clin Exper Allergy. 2011; 42: 30-7.
- 8. Fontaine M, et al. [Severe bronchospasm using Diprivan® in a patient allergic to peanut and birch]. Ann Fr Anesth Reanim. 2011 Feb;30(2):147-9.{PubMed abstract]
- 9. Canadian Food Insepction Agency. Sesame allergy. Available from URL: http://www.inspection.gc.ca/english/fssa/labeti/allerg/sese.shtml. Accessed 15/3/2012.
- 10. Lunn M, Fausnight T. Hypersensitivity to total parenteral nutrition fat-emulsion component in an egg-allergic child. Pediatrics. 2011 Oct;128(4):e1025-8. (see also Murphy et al. Anesth Analg. 2011; 113: 140-4.)
- 11. Canadian Immunization Guide, 7th ed. (2006). Available from URL: http://www.phac-aspc.gc.ca/publicat/cig-gci/. Accessed 15/3/2012.
- 12. National Advisory Committee on Immunization. Statement on seasonal influenza vaccine for 2011-2012. Available from URL: http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/11vol37/acs-dcc-5/index-eng.php. Accessed 15/3/2012.
- 13. Sapone A, et al. Spectrum of gluten-related disorders: consensus on new nomenclature and classification. BMC Med. 2012; 10: 13 (12 pages).
- 14. Mukadam ME, et al. Management during cardiopulmonary bypass of patients with

- presumed fish allergy. J Cardiothor Vasc Anesth. 2001; 15: 512-9. (See also Knape JTA, et al. Anethesiol. 1981; 55: 324-5 and Dewachter P, et al. Curr Opin Anethesiol. 2011; 24: 320-5.)
- 15. Katelaris CH. 'lodine allergy' label is misleading. Aust Prescr. 2009; 32:125-8.
- 16. Bukacel DG, et al. Cross-reactivity between paclitaxel and hazelnut: a case report. J Oncol Pharm Pract. 2007; 13: 53-5.
- 17. Cunningham E. Are there foods that should be avoided if a patient is sensitive to salicylates? J Am Diet Assoc. 2010; 110: 976.
- 18. Greenhawt M, et al. Carmine sensitivity masquerading as azithromycin hypersensitivity. Allergy Asthma Proc. 2009; 30: 95-101.
- 19. Salicylates general statement. AHFS DI (online via MedicinesComplete)
- Ozdemir O, et al. Development of multiple food allergies in children taking tacrolimus after heart and liver transplantation. Pediatr Transplant. 2006; 10: 380-3. (See also: Yilmaz et al. Transplant Proc. 2005; 37: 4251-3 and Martin-Munoz MF, et al. Br J Dermatol. 2005; 153: 842-3.)
- 21. Vickery P, et al. Pathophysiology of food allergy. Pediatr Clin N Am. 2011; 363-76.
- 22. Osborn DA, Sinn JK. Probiotics in infants for prevention of allergic disease and food hypersensitivity. Cochrane Database Syst Rev. 2007 Oct 17;(4):CD006475.
- 23. Personal communication: epinephrine dosing, information on file (DPIC)
- 24. http://www.allergysafecommunities.ca

©2012 B.C. Drug and Poison Information Centre

A version of this document was published in BCPhA's The Tablet. 2012; 20(8): 8-9.

We are grateful to all the First Nations who have cared for and nurtured the lands and waters around us for all time, including the x?m??k??y??m (Musqueam), Sk?wx?wu?7mesh U?xwumixw (Squamish Nation), and s?l?ílw?ta? (Tsleil-Waututh Nation) on whose unceded and ancestral territory our centre is located.

© 2024 BC Drug and Poison Information Centre

All material found on the BC Drug and Poison Information Centre (DPIC) website is provided for informational purposes only. It is *not* meant to replace the expert advice of a healthcare professional such as a physician, pharmacist, nurse or qualified poison specialist. Use of this site is governed and restricted by specific terms of use. Please review the **full terms and conditions** below prior to using the DPIC website. In the event of a poisoning emergency, call your local poison control centre immediately. Portions of this web site are intended for healthcare professionals. Interpretation and application of information may require more detailed explanation than contained herein, particularly regarding any clinical information that is found in or linked to this site. Patients are advised to consult their health care provider regarding diagnosis and treatment, and for assistance in interpreting these materials and applying them in individual cases.

Terms and Conditions