



Published on *British Columbia Drug and Poison Information Centre (BC DPIC)* (<http://www.dpic.org>)

[Home](#) > [Printer-friendly PDF](#) > [Printer-friendly PDF](#)

---

## Niacin: The facts on flushing

### Access:

professional

### Article type:

drug information

The BC Drug & Poison Information Centre commonly receives calls from patients who are experiencing an adverse reaction to over-the-counter niacin. Typically patients are taking the drug for the first time in higher doses for hypercholesterolemia or other indications. Usually, they are quite concerned about an "allergic" reaction to the vitamin. Here are some commonly asked questions and answers about flushing from niacin:

**Q: I recently started taking niacin for high cholesterol. The first time I took it I experienced a hot flushing sensation on my skin that took an hour to go away. I thought I was having an allergic reaction. Can you tell me what happened?**

*Niacin (nicotinic acid vitamin)*, or vitamin B<sub>3</sub>, is an essential vitamin. Adults require a daily dietary intake of about 15 mg. At much higher doses (1000 to 2000 mg per day) niacin is used as a treatment for high cholesterol.

These higher doses of niacin cause an intense flushing or "prickly heat" sensation to the face and upper body, usually 15-30 minutes after taking a relatively large dose (e.g. 500 mg). This flushing is experienced by almost everyone and, while it might feel like an allergy, it is not a true allergic reaction.

Other than causing discomfort the flushing is harmless and usually subsides within 1 or 2 hours. Symptoms are most intense after the first dose and typically diminish over days or weeks with continued use of niacin.

**Q: Are there different types of niacin that might reduce the flushing?**

The flushing occurs with over-the-counter immediate-release niacin tablets. With this type of niacin the vitamin is delivered to the body in a short burst and the flushing reaction is more intense.

Sustained-release niacin tablets deliver the vitamin to the body in a slower fashion over many hours. This reduces the intensity of the flushing but this type of niacin causes liver damage in some people.

A prescription extended-release niacin product called Niaspan® releases niacin in a slower way but over a shorter period of time compared to the sustained-release tablets. This gives the liver a "break" from processing the niacin making liver damage less likely. This type of niacin has been shown to have positive effects on cholesterol with reduced flushing. However, it is more expensive than regular niacin tablets.

**Q: What about "no-flush" niacin?**

No-flush niacin contains something called inositol nicotinate, which the body is supposed to slowly convert to niacin. However, there is evidence that it does not actually provide the body with much niacin. This is probably the reason it does not produce any flushing. Of course, this also means that no-flush niacin does not have any of the beneficial effects on cholesterol.

**Q: If I continue to take the regular niacin tablets, are there any tips to minimize the flushing?**

The key to reducing the intensity of niacin flushing is to start with a low dose and gradually increase the dose over a period of weeks. Taking it with food also helps reduce the intensity of the reaction.

One approach is to start immediate-release niacin at 100 mg twice daily after a meal for the first week, then double the daily dose each week until you are taking what the doctor has prescribed.

Aspirin will also help to reduce the flushing. If you are already taking low-dose aspirin (81 to 325 mg daily) try taking it about 30 minutes before your first niacin dose of the day.

**References:**

1. Anon. Don't overlook niacin for treating cholesterol problems. Harvard Heart Letter. April

2004.

2. Jacobson TA. A "hot" topic in dyslipidemia management - "How to beat a flush": Optimizing niacin tolerability to promote long-term treatment adherence and coronary disease prevention. *Mayo Clin Proc.* 2010;85(4):365-79.
3. Lee T. Is no-flush niacin as effective as other kinds of niacin? *Harvard Heart Letter.* March 2010.
4. Meyers CD, et al. Varying cost and free nicotinic acid content in over-the-counter niacin preparations for dyslipidemia. *Ann Intern Med.* 2003;139(12):996-1002.

by Rob Gair, BSc (Pharm), CSPI

### ©2013 B.C. Drug and Poison Information Centre

*A version of this document was published in BCPHA's The Tablet. 2013; 22(2): 10-11.*

**Keywords:** niacin  
flushing

© 2017 BC Drug and Poison Information Centre

All material found on the BC Drug and Poison Information Centre (DPIC) website is provided for informational purposes only. It is *not* meant to replace the expert advice of a healthcare professional such as a physician, pharmacist, nurse or qualified poison specialist. Use of this site is governed and restricted by specific terms of use. Please review the **full terms and conditions** below prior to using the DPIC website. In the event of a poisoning emergency, call your local poison control centre immediately. Portions of this web site are intended for healthcare professionals. Interpretation and application of information may require more detailed explanation than contained herein, particularly regarding any clinical information that is found in or linked to this site. Patients are advised to consult their health care provider regarding diagnosis and treatment, and for assistance in interpreting these materials and applying them in individual cases.

#### **Terms and Conditions**

---

**Source URL (retrieved on 2019-10-20 18:24):** <http://www.dpic.org/article/professional/niacin-facts-flushing>